

**Hematologic and Serum Biochemical Data from
Mass Stranded Long-Finned Pilot Whales
(*Globicephala melas*), Cape Cod, USA, 2002**

Supplemental Table 1

Garrett C. Crooks,¹ Sarah M. Sharp,² Constance Merigo,³
Kathleen M. Moore,² and Charles J. Innis³

¹Colorado State University, 300 W. Drake, Fort Collins, CO 80523, USA
E-mail: Garrett.Crooks@colostate.edu

²International Fund for Animal Welfare, 290 Summer Street, Yarmouth Port, MA 02675, USA

³New England Aquarium, 1 Central Wharf, Boston, MA 02110, USA

Pilot Whale Hematology and Chemistry

| | | | | | | | | | | | | | | | | | | | | |
|----------------------------|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Globulin (g/dL) | NE | 4 | 4.1 | 3 | 5.3 | 2.8 | 3.8 | 3.3 | 3.2 | 4 | 3.6 | 3.9 | 3.5 | 2.2 | 3.8 | 2.5 | 3.7 | 3.9 | 2.7 | 3.2 |
| A/G ratio | NE | 0.9 | 0.8 | 1.2 | 0.8 | 1.2 | 1.0 | 1.1 | 1.1 | 0.8 | 1.0 | 0.9 | 0.9 | 1.6 | 1.1 | 1.5 | 1.0 | 1.0 | 1.6 | 1.2 |
| α 1 globulin (g/dL) | NE | 0.5 | 1.0 | 0.5 | 0.7 | 0.7 | 0.5 | 0.7 | 0.4 | 0.5 | 0.5 | 0.4 | 0.5 | 0.9 | 0.8 | 1.0 | 0.5 | 0.5 | 0.8 | 0.4 |
| α 2 globulin (g/dL) | NE | 0.8 | 0.8 | 0.6 | 0.8 | 0.5 | 0.7 | 0.9 | 0.8 | 0.7 | 0.8 | 0.7 | 0.7 | 0.4 | 0.6 | 0.4 | 0.7 | 0.9 | 0.2 | 0.8 |
| β 1 globulin (g/dL) | NE | 0.2 | 0.4 | 0.2 | 0.6 | 0.2 | 0.3 | 0.4 | 0.4 | 0.2 | 0.8 | 0.2 | 0.2 | 0.3 | 0.5 | 0.6 | 0.2 | 0.5 | 0.2 | 0.4 |
| β 2 globulin (g/dL) | NE | 0.2 | ND | ND | ND | 0.3 | ND | ND | ND | 0.3 | ND | 0.3 | 0.3 | ND | ND | ND | 0.3 | ND | ND | ND |
| γ 1 globulin (g/dL) | NE | 2.3 | 2.1 | 1.8 | 3.3 | 1.2 | 2.4 | 1.3 | 1.5 | 2.3 | 1.6 | 2.4 | 1.8 | 0.7 | 1.9 | 0.6 | 2.0 | 1.6 | 1.5 | 1.6 |