

Short Note

Movements and Timing of Humpback Whales (*Megaptera novaeangliae*) Within the Breeding Region of the Eastern South Pacific

Camila A. Valdivia,¹ Aldo S. Pacheco,² Fernando Félix,^{3,4} Ben Haase,³
Kristin Rasmussen,⁵ Luis Santillán,^{6,7} Belen Alcorta,⁸ and Sebastian Silva⁸

¹*Facultad de Ciencias del Mar y Recursos Biológicos, Universidad de Antofagasta,
Avenida Universidad de Antofagasta 02800, Antofagasta, Chile*

²*Instituto de Ciencias Naturales Alexander von Humboldt, CENSOR Laboratory, Universidad de Antofagasta,
Avenida Universidad de Antofagasta 02800, Antofagasta, Chile
E-mail: babuchapv@yahoo.com*

³*Museo de Ballenas, Avenida Enríquez Gallo S/N, Salinas, Ecuador*

⁴*Pontificia Universidad Católica del Ecuador (PUCE), Facultad de Ciencias Exactas y Naturales,
Avenida 12 de Octubre y Roca, Quito, Ecuador*

⁵*Panacetacea, David, Chiriquí, Panamá*

⁶*Universidad San Ignacio de Loyola, Perú, Avenida La Fontana 550, Lima 12, Perú*

⁷*Centro Peruano de Estudios Cetológicos (CEPEC), Museo de los Delfines, Pucusana, Lima 20, Perú*

⁸*Pacífico Adventures, Avenida Rivera del Mar s/n, Los Organos, Perú*

Eastern South Pacific humpback whales (*Megaptera novaeangliae*) (Breeding Stock G, according to the International Whaling Commission [IWC], 1998) migrate from feeding grounds located mainly in the surroundings of the Antarctic Peninsula (Dalla Rosa et al., 2008) and to a lesser extent in the Magellan Strait (Gibbons et al., 2003; Acevedo et al., 2006, 2013) and Golfo de Corcovado in southern Chile (Hucke-Gaete et al., 2013) to their breeding region off northern Peru, Ecuador, Colombia, Panama, and Costa Rica (Flórez-González, 1991; Flórez-González et al., 1998; Félix & Haase, 2001; Rasmussen et al., 2007; Pacheco et al., 2009; Guidino et al., 2014; Guzmán et al., 2015). The connectivity between some feeding and breeding grounds has been demonstrated through photo-identification (Stone et al., 1990; Stevick et al., 2004; Acevedo et al., 2007; Rasmussen et al., 2007; Capella et al., 2008) and to a lesser extent with genetics (Caballero et al., 2001; Félix et al., 2012). In addition, regional movements of five individuals and a migratory path used by one humpback whale from the breeding region to feeding grounds has been recently characterized via satellite tracking (Félix & Guzmán, 2014). Although important information on connectivity and migration at the geographical scale has been generated, little is known about the movement patterns at the regional or mesoscale.

Humpback whales tend to have a continuous distribution throughout coastal areas of the breeding region (Zerbini et al., 2004; Félix & Haase, 2005; Van Waerebeek et al., 2013). However, residence patterns and movements within the breeding region may vary within and between seasons (Baker & Herman, 1981; Calambokidis et al., 2001; Smith et al., 2012; Guzmán et al., 2015). Understanding the movement patterns within the breeding area may provide valuable information about the migration timing and habitat use by whales along the breeding grounds (e.g., Guzmán & Félix, 2017). This information may help to identify potential risks to humpback whales caused by human activities such as interaction with gill-nets (García-Godos et al., 2013), and seismic and whale-watching operations (Pacheco et al., 2011); and it may provide the understanding to address them.

Previous photo-identification studies in the eastern South Pacific revealed the connectivity between sites within the breeding region—Ecuador, Colombia, Panama, and Costa Rica (Flórez-González et al., 1998; Castro et al., 2008; Félix et al., 2009). Northern Peru was considered in those studies, but the sample size was rather low (no more than 20 photographs). Research efforts in the region have increased progressively in the last 20 y (Flórez-González et al., 2007), and more sites within the breeding region have

been surveyed (Pacheco et al., 2009; Castro et al., 2011; Rojas et al., 2014). These studies support the notion that this population has a continuous distribution along the eastern South Pacific from northern Peru to southern Costa Rica during the austral winter/spring breeding season.

Information on variability in abundance may provide some insights about timing and movements of eastern South Pacific humpback whales within the breeding region. Thus, an apparent relationship would exist between whale abundance and latitude. Whale abundance peaks off Panama in August and September (Guzmán et al., 2015) and off Colombia in August (Flórez-González et al., 1998); while off Ecuador, it is reported in July (Scheidat et al., 2000; Félix & Haase, 2001). In northern Peru, the peak is observed towards the end of the breeding season in September and October (Pacheco et al., 2009; Santillán, 2011; Guidino et al., 2014). This variability in abundance suggests that whales have an offshore path arriving first to Ecuador in July, then moving up north to Colombia, and finally moving further north to Panama. The peak of abundance later in the season (September and October) in northern Peru would indicate that this southern region constitutes a migratory corridor for humpback whales returning to feeding areas via neritic waters. This belief is also supported by whaling statistics from Paita (*ca.* 5° S), northern Peru, which show the landing peak for humpback whales in October (Ramírez, 1988).

Herein, we tested the prediction that humpback whales would primarily use northern destinations early in the breeding season—first to Ecuador and then to Panama, and then move towards the south via northern Peru. To study this prediction, we compared photo-identified humpback whales (*i.e.*, photographs of the ventral side of the flukes [see Katona et al., 1979] and also the dorsal fin in some cases) coming from the neritic zones of northern Peru with those of central Ecuador and Panama, thus covering the major extension of the breeding region (Figure 1).

Photographs were taken between June and October during the 1991 to 2014 study period, principally during whale-watching excursions but also during studies of humpback whale distribution and population dynamics (Table 1). For the comparison, photographs were simultaneously visualized using a double screen and examined by the naked eye. Photographs with poor resolution, bad quality, and duplicates were discarded. Catalogs were compared in the following sequence: Peru vs Ecuador, Peru vs Panama, and Ecuador vs Panama.

A total of 2,999 unique photo-identified humpback whales were compared yielding 39

matches (19 photographs were discarded due to poor resolution; see Table 1). Only five matches were detected within the same season. In 2010, two individuals sighted in Ecuador in July were resighted in northern Peru—one in August and one in October. A third animal was sighted off Ecuador on 8 September and seen 4 d later off northern Peru (Table 2). (Photographs of the total matches are available on the *Aquatic Mammals* website: www.aquaticmammalsjournal.org/index.php?option=com_content&view=article&id=10&Itemid=147). The estimated time elapsed between locations was 34.9 d (SD = 21.2) on average. Another within-season resighting was made in 2008: an individual sighted in Panama in July was resighted in Ecuador in August. Finally, an individual sighted in Panama in August 2013 was resighted in northern Peru in September (Table 2). The estimated time between locations was 45 and 36 d, respectively. Of the 34 between-years matches, 30 were between Ecuador and Peru, and four were between Panama and Ecuador (Table 2). The timing of sighting and resighting in these cases was similar as in the within-seasons analysis. Individuals were sighted first in Ecuador and/or Panama during June, July, and August, and then whales were resighted in northern Peru after variable times ranging from 2 wks to 3 mo during August, September, and October (Table 2). The longest estimated time interval was between 17 July to 30 October. Only five resightings showed the opposite timing (Table 2). A chi square test comparing the timing of recaptures between Peru and Ecuador provided statistical significance to the observed tendency of sighting whales early in the season in Ecuador and late in Peru ($\chi^2 = 29.9$; $df = 4$, $p < 0.05$). No statistical comparison was made with Panama due to the low number of recaptures.

Although humpback whales arrive first to Ecuador and then to Panama, matches indicated that most individuals moved in the opposite direction (south) as the season progresses. Persistent and directional movement seems to be a common characteristic for humpback whales in breeding areas. For example, Baker & Herman (1981) reported a southeast to northwest movement pattern in the breeding area at the Hawaiian Archipelago. Smith et al. (2012) modeled the habitat used by breeding humpback whales in eastern Australia and suggested that whales arrive first to neritic waters at northern breeding destinations and then move southwards closer to the coast. Indeed, southern locations were considered migratory corridors.

A plausible explanation of such distribution within breeding grounds could be a consequence of population stratification, with humpback whales arriving first to the breeding destination,




Figure 1. The map on the right shows the breeding region for humpback whales (*Megaptera novaeangliae*) of the eastern South Pacific. The map on the left indicates the locations (open dots) where photographic surveys were conducted. The grey arrow highlights the main north to south movement revealed in this study (see “Results” for details).

Table 1. Number of humpback whales (*Megaptera novaeangliae*) identified by photographs of flukes at the different study sites along the breeding region

Institutions	No. of photographs		Years	Sightings*	Observation time†	Location	Country
	Available	Examined					
Panacetacea	445	441	2002-2013	946	1,188	Golfo de Chiriqui	Panamá
MBS	222	218	1991-1997	180	94	Puerto López	Ecuador
MBS	1,898	1,890	2001-2013	1,907	855	1° 24' S, 80° 55' W Salinas	Ecuador
Pacífico Adventures	440	437	2009-2014	1,602	372	2° 13' S, 80° 57' W Los Organos	Perú
CEPEC	13	13	2009-2010	43	82	4° 10' S, 81° 8' W Sechura	Perú
Total	3,018	2,999		4,678	2,591	5° 34' S, 80° 57' W	

CEPEC = Centro Peruano de Estudios Cetológicos, MBS = Museo de Ballenas de Salinas, * = total number of sighted whales, and † = total number of observation hours.

Table 2. Photograph matches among humpback whales from Peru, Ecuador, and Panama along the eastern South Pacific breeding region. The numbers in the first column are the time intervals between matches in days; the first five rows are within-year recaptures followed by the rest of the between-years matches.

	Perú				Ecuador		Panamá	
	Sechura		Los Organos		Salinas/Puerto López		Chiriqui	
Time	ID	Date	ID	Date	ID	Date	ID	Date
20			PAO#0026	8 Aug. 2010	MBS#1687	18 July 2010		
103	SE 09-4	30 Oct. 2010			MBS#1870	17 July 2010		
4			PAO#0004	12 Sept. 2010	MBS#1949	8 Sept. 2010		
45					MBS#1414	30 Aug. 2008	PAN#1106	15 July 2008
36			PAO#0305	27 Sept. 2013			PAN#1321	21 Aug. 2013
6			PAO#0392	16 Sept. 2014	MBS#0116	22 Sept. 1996		
8			PAO#0106	11 Aug. 2012	MBS#0748	3 Aug. 2010		
7					MBS#2104	3 Aug. 2012	PAN#1182	10 Aug. 2007
9			PAO#0013	29 Aug. 2009	MBS#0059	20 Aug. 1995		
13			PAO#0004	15 Aug. 2009	MBS#2009	2 Aug. 2008		
14					MBS#1527	31 July 2009	PAN#1076	14 Aug. 2007
14			PAO#0031	14 Aug. 2010	MBS#0771	31 July 2006		
15			PAO#0164	19 Sept. 2012	MBS#1440	4 Sept. 2008		
17			PAO#0127	27 Aug. 2012	MBS#0181	10 Aug. 2010		
49			PAO#0050	5 Sept. 2010	MBS#0988	16 July 2006		
19			PAO#0023	5 Aug. 2010	MBS#0739	16 July 2006		
20			PAO#0282	18 Sept. 2013	MBS#0947	29 Aug. 2006		
21			PAO#0109	15 Aug. 2012	MBS#0391	6 Sept. 2003		
26			PAO#0051	14 Sept. 2010	MBS#1155	19 Aug. 2007		
27			PAO#0074	26 Aug. 2011	MBS#0948	29 July 2006		
29			PAO#0010	9 Aug. 2009	MBS#0253	8 Sept. 2008		
66			PAO#0279	13 Sept. 2013	MBS#0517	7 July 2004		
37			PAO#0014	14 Sept. 2009	MBS#0453	7 Aug. 2004		
38			PAO#0314	30 Sept. 2013	MBS#1904	22 Aug. 2010		
42			PAO#0015	14 Sept. 2009	MBS#0592	2 Aug. 2005		
42			PAO#0318	13 Oct. 2013	MBS#1193	1 Sept. 2007		
44			PAO#0055	19 Sept. 2010	MBS#1095	5 Aug. 2008		
44			PAO#0059	30 Sept. 2010	MBS#0011	16 Aug. 1992		
44			PAO#0077	24 Aug. 2011	MBS#1488	10 July 2009		
45			PAO#0006	21 Aug. 2009	MBS#0255	6 July 2002		
47			PAO#0245	19 Sept. 2010	MBS#1270	2 Aug. 2007		
48			PAO#0056	19 Aug. 2010	MBS#1155	1 July 2009		
48			PAO#0257	3 Sept. 2013	MBS#1683	15 July 2008		
52					MBS#1630	14 Sept. 2010	PAN#1287	22 July 2007
53			PAO#0009	10 Sept. 2013	MBS#1002	17 July 2010		
57					MBS#1658	11 July 2010	PAN#1365	8 Sept. 2013
60			PAO#0173	22 Sept. 2012	MBS#0752	22 July 2006		
63			PAO#0323	12 Oct. 2013	MBS#0372	9 Aug. 2003		
66	SE09-10	30 Oct. 2010			MBS#0168	24 Aug. 1997		

following an offshore path, and then being resighted in the southern areas by the end of the season (September-October) once the southern coastal migration has started. This change to a coastal distribution is more evident in females with calves (Félix & Botero-Acosta, 2011;

Guidino et al., 2014), a behavior that is maintained by migrating females after leaving the breeding area (Félix & Guzmán, 2014). Recently generated information on movements of whales that were satellite-tagged in Panama and Ecuador supports stratification in this population at breeding

grounds (Guzmán & Félix, 2017) as well as the low number of matches in a large sample as in our study.

We are aware that migration is a complex process with different timing and overlapping stages (Dawbin, 1966; Craig et al., 2003), and spatial segregation according to the whale's age/class (Félix & Haase, 2005; Félix & Botero-Acosta, 2011; Rasmussen et al., 2012; Guidino et al., 2014), which increases uncertainty and precludes reaching definitive conclusions from the available data. In the same line, we suggest caution with between-years matching results. Although these results point to a consistent movement pattern, we cannot unequivocally account for the explanatory processes behind this since whale behavior may also vary between years due to environmental variability. Other factors must also be considered such as short residence time and the relatively large home range (between 25,000 and 60,00 km²; *sensu* Guzmán & Félix, 2017) which may not be covered entirely during photographic surveys. In addition, it may be difficult to distinguish between the absence of an individual in a group and the failure to capture it photographically, and the low probability of capture of fluke-up which is lower in breeding areas with respect to feeding areas. Moreover, the photographic effort may vary throughout the breeding area as research teams do not necessarily survey humpback whales simultaneously across their study region during the whole breeding season. Nevertheless, our comparison and previous studies (e.g., Castro et al., 2008; Félix et al., 2009) call attention to this north-to-south movement pattern because of the implications in population estimate studies using mark-recapture models since such heterogeneity may affect whales' probability of capture (Félix et al., 2011). This study highlights the importance of monitoring movements of humpback whales within the breeding area of the eastern South Pacific as in other regions. In future studies, we recommend including photographs from Colombia to cover the entire breeding area as well as to increase telemetry studies, which would enhance our understanding of humpback whale movements during the breeding season.

Acknowledgments

C. Valdivia was supported by the following scholarship: *Apoyo a las actividades de la titulación de la Vicerectoría de Investigación Innovación y Post-grado, Universidad de Antofagasta*. K. Rasmussen was supported by The Moore Charitable Foundation and the Islas Secas Resort. L. Santillán received support from the Rufford Small Grant Program and Prodelphinus NGO for

sampling at Sechura. A. S. Pacheco, S. Silva, and B. Alcorta sincerely thank the crew and volunteers of Pacifico Adventures for their enormous contribution on the collection of photographs at Los Organos. Comments by two anonymous reviewers and Manuel dos Santos helped us to improve an earlier version of this manuscript.

Literature Cited

- Acevedo, J., Aguayo-Lobo, A., & Pastene L. A. (2006). Filopatría de la ballena jorobada (*Megaptera novaeangliae* Borowski, 1781), al área de alimentación del estrecho de Magallanes [Site fidelity of humpback (*Megaptera novaeangliae* Borowski, 1781) to the area of the Strait of Magellan]. *Revista de Biología Marina y Oceanografía*, 41(1), 11-19. <https://doi.org/10.4067/S0718-19572006000100004>
- Acevedo, J., Plana, J., Aguayo-Lobo, A., & Pastene, L. A. (2011). Surface feeding behavior of humpback whales in the Magellan Strait. *Revista de Biología Marina y Oceanografía*, 46(3), 483-490. <https://doi.org/10.4067/S0718-19572011000300018>
- Acevedo, J., Haro, D., Dalla-Rosa, L., Aguayo-Lobo, A., Huckle-Gaete, R., Secchi, E., . . . Pastene, L. A. (2013). Evidence of spatial structuring of eastern South Pacific humpback whale feeding grounds. *Endangered Species Research*, 22(1), 33-38. <https://doi.org/10.3354/esr00536>
- Acevedo, J., Rasmussen, K., Félix, F., Castro, C., Llano, M., Secchi, E., . . . Pastene, L. A. (2007). Migratory destinations of the humpback whales from Magellan Strait feeding ground, Chile. *Marine Mammal Science*, 23(2), 453-463. <https://doi.org/10.1111/j.1748-7692.2007.00116.x>
- Baker, C. S., & Herman, L. M. (1981). Migration and local movement of humpback whales (*Megaptera novaeangliae*) through Hawaiian waters. *Canadian Journal of Zoology*, 59(3), 460-469. <https://doi.org/10.1139/z81-067>
- Caballero, S., Hamilton, H., Jaramillo, H., Capella, J., Flórez-González, L., Olavarría, C., . . . Baker, C. S. (2001). Genetic characterisation of the Colombian Pacific Coast humpback whale population using RAPD and mitochondrial DNA sequences. *Memoirs of the Queensland Museum*, 47(2), 459-464.
- Calambokidis, J., Steiger, G. H., Straley, J. M., Herman, L. M., Cerchio, S., Salden, D. R., . . . Quinn, T. J. (2001). Movements and population structure of humpback whales in the North Pacific. *Marine Mammal Science*, 17(4), 769-794. <https://doi.org/10.1111/j.1748-7692.2001.tb01298.x>
- Capella, J. J., Gibbons, J., Flórez-González, L., Llano, M., Valladares, C., Sabaj, V., & Vilina, Y. A. (2008). Migratory round-trip of individually identified humpback whales at the Strait of Magellan: Clues on transit times and phylopatry to destinations. *Revista Chilena de Historia Natural*, 81(4), 547-560. <https://doi.org/10.4067/S0716-078X2008000400008>
- Castro, C., Acevedo, J., Allen, J., Dalla Rosa, L., Flórez-González, L., Aguayo-Lobo, A., . . . Pastene, L. A.

- (2008, June). *Migratory movements of humpback whales (Megaptera novaeangliae) between Machalilla National Park, Ecuador and southeast Pacific* (Paper SC/60/SH). Presented at the 60th Annual Meeting of the Scientific Committee of the International Whaling Commission. Santiago de Chile, Chile.
- Castro, C., Alcorta, B., Allen, J., Cáceres, C., Forestell, P., Kauffman, G., . . . Tagarino, A. (2011, June). *Comparison of the humpback whale catalogues between Ecuador, Peru and American Samoa evidence of the enlargement of the Breeding Stock G to Peru* (Paper SC/63/SH19). Presented at the 63rd Annual Meeting of the Scientific Committee of the International Whaling Commission. Tromsø, Norway.
- Craig, A. S., Herman, L. M., Gabriele, C. M., & Pack, A. A. (2003). Migratory timing of humpback whales (*Megaptera novaeangliae*) in the central North Pacific varies with age, sex and reproductive status. *Behavior*, *140*, 981-1001. <https://doi.org/10.1163/156853903322589605>
- Dalla Rosa, L., Secchi, E. R., Maia, Y. G., Zerbini, A. N., & Heide-Jørgensen, M. P. (2008). Movements of satellite-monitored humpback whales on their feeding ground along the Antarctic Peninsula. *Polar Biology*, *31*(7), 771-781. <https://doi.org/10.1007/s00300-008-0415-2>
- Dawbin, W. H. (1966). The seasonal migratory cycle of humpback whales. In K. S. Norris (Ed.), *Whales, dolphins, and porpoises* (pp. 145-170). Berkeley: University of California Press. xv + 789 pp.
- Félix, F., & Botero-Acosta, N. (2011). Distribution and behaviour of humpback whale mother-calf pairs during the breeding season off Ecuador. *Marine Ecology Progress Series*, *426*, 277-287. <https://doi.org/10.3354/meps08984>
- Félix, F., & Guzmán, H. (2014). Satellite tracking and sighting data analyses of southeast Pacific humpback whales (*Megaptera novaeangliae*): Is the migratory route coastal or oceanic? *Aquatic Mammals*, *40*(4), 329-340. <https://doi.org/10.1578/AM.40.4.2014.329>
- Félix, F., & Haase, B. (2001). The humpback whale off the coast of Ecuador, population parameters and behavior. *Revista de Biología Marina y Oceanografía*, *36*(1), 61-74. <https://doi.org/10.4067/S0718-19572001000100006>
- Félix, F., & Haase, B. (2005). Distribution of humpback whales along the coast of Ecuador and management implications. *Journal of Cetacean Research and Management*, *7*(1), 21-31.
- Félix, F., Caballero, S., & Olavarría, C. (2012). Genetic diversity and population structure of humpback whales (*Megaptera novaeangliae*) from Ecuador based on mitochondrial DNA analyses. *Journal of Cetacean Research and Management*, *12*(1), 71-77.
- Félix, F., Castro, C., Laake, J., Haase, B., & Scheidat, M. (2011). Abundance and survival estimates of the southeastern Pacific humpback whale stock from 1991-2006 photo-identification surveys in Ecuador. *Journal of Cetacean Research and Management*, *3*(Special Issue), 301-307.
- Félix, F., Rasmussen, K., Garita, F., Haase, B., & Simonis, A. (2009, June). *Movements of humpback whales between Ecuador and Central America, wintering area of the breeding stock G* (Paper SC/61/SH18). Presented at the 61st Annual Meeting of the Scientific Committee of the International Whaling Commission. Madeira, Portugal.
- Flórez-González, L. (1991). Humpback whales *Megaptera novaeangliae* in the Gorgona Island, Colombian Pacific breeding waters: Population and pod characteristics. *Memoirs of the Queensland Museum*, *30*(2), 291-295.
- Flórez-González, L., Capella, A. J., Haase, B., Bravo, G. A., Félix, F., & Gerrodette, T. (1998). Changes in winter destinations and the northernmost record of southeastern Pacific humpback whales. *Marine Mammal Science*, *14*(1), 189-196. <https://doi.org/10.1111/j.1748-7692.1998.tb00707.x>
- Flórez-González, L., Ávila, I. C., Alzueta, J. C., Falk, P., Félix, F., Gibbons, J., . . . Van Waerebeek, K. (2007). *Estrategia para la conservación de la ballena jorobada del Pacífico sudeste: Lineamientos para un plan de acción regional e iniciativas nacionales* [Conservation strategy for southeast Pacific humpback whale: Guidelines for a regional action plan and national initiatives]. Cali, Colombia: Fundación Yubarta. 106 pp.
- García-Godos, I., Van Waerebeek, K., Alfaro-Shigueto, J., & Mangel, J. (2013). Entanglements of large cetaceans in Peru: Few records but high risk. *Pacific Science*, *67*(4), 523-532. <https://doi.org/10.2984/67.4.3>
- Gibbons, J., Capella, J. C., & Valladares, C. (2003). Rediscovery of a humpback whale (*Megaptera novaeangliae*) feeding ground in the Straits of Magellan, Chile. *Journal of Cetacean Research and Management*, *5*(2), 203-208.
- Guidino, C., Llapapasca, M., Silva, S., Alcorta, B., & Pacheco, A. S. (2014). Patterns of spatial and temporal distribution of humpback whales at the southern limit of the southeast Pacific breeding area. *PLOS ONE*, *9*(11), e112627. <https://doi.org/10.1371/journal.pone.0112627>
- Guzmán, H., & Félix, F. (2017). Movements and habitat use by southeast Pacific humpback whales satellite tracked at two breeding sites. *Aquatic Mammals*, *43*(2), 139-155. <https://doi.org/10.1578/AM.43.2.2017.139>
- Guzmán, H. M., Condit, R., Pérez-Ortega, B., Capella, J. J., & Stevick, P. T. (2015). Population size and migratory connectivity of humpback whales wintering in Las Perlas Archipelago, Panama. *Marine Mammal Science*, *31*(1), 90-105. <https://doi.org/10.1111/mms.12136>
- Hucke-Gaete, R., Haro, D., Torres-Flóres, J. P., Montecinos, J., Viddi, F., Bedriñana-Romano, L., . . . Ruiz, J. (2013). A historical feeding ground for humpback whales in the eastern South Pacific revisited: The case of northern Patagonia, Chile. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *23*, 858-867. <https://doi.org/10.1002/aqc.2343>
- International Whaling Commission (IWC). (1998). Report of the Scientific Committee, Annex G. Report of the Sub-Committee on the Comprehensive Assessment of Southern Hemisphere Humpback Whales. *Report of*

- the International Whaling Commission*, 48, 170-182. Retrieved from <https://iwc.int/annual-reports-pre-1998>
- Katona, S., Baxter, B., Brazier, O., Kraus, S., Perkins, J., & Whitehead, H. (1979). Identification of humpback whales by fluke photographs. In H. E. Winn & B. L. Olla (Eds.), *Behavior of marine animals* (Chapter 2, pp. 33-34). New York: Plenum Press. https://doi.org/10.1007/978-1-4684-2985-5_2
- Pacheco A. S., Silva, S., & Alcorta, B. (2009). Winter distribution and group composition of humpback whales (*Megaptera novaeangliae*) off northern Peru. *Latin American Journal of Aquatic Mammals*, 7(1-2), 33-38. <https://doi.org/10.5597/lajam00131>
- Pacheco A. S., Silva, S., & Alcorta, B. (2011). Is it possible to go whale-watching in the coast off Peru? A study case with humpback whales. *Latin American Journal of Aquatic Research*, 39(1), 189-196. <https://doi.org/10.3856/vol39-issue1-fulltext-20>
- Ramírez, P. (1988). La ballena jorobada *Megaptera novaeangliae* en la costa Norte del Perú [The humpback whale *Megaptera novaeangliae* on the northern coast of Peru]. *Boletín de Lima*, 56, 91-96.
- Rasmussen, K., Calambokidis, J., & Steiger, G. H. (2012). Distribution and migratory destinations of humpback whales off the Pacific coast of Central America during the boreal winters of 1996-2003. *Marine Mammal Science*, 28(3), E267-E279. <https://doi.org/10.1111/j.1748-7692.2011.00529.x>
- Rasmussen, K., Palacios, D. M., Calambokidis, J., Saborío, M. T., Rosa, L. D., Secchi, E. R., . . . Stone, G. S. (2007). Southern hemisphere humpback whales wintering off Central America: Insights from water temperature into the longest mammalian migration. *Biology Letters*, 3(3), 302-330. <https://doi.org/10.1098/rsbl.2007.0067>
- Rojas, K., Denkinger, J., Arahana, V., Dalgo, D., & Torres, M. D. L. (2014). Analysis of the site fidelity between male and female humpback whales, visiting the Esmeraldas coasts (Ecuador). *Avances en Ciencias e Ingenierías*, 6(1), B28-B35.
- Santillán, L. (2011). Records of humpback whales (*Megaptera novaeangliae*) in Sechura Bay, Peru, in spring 2009-2010. *Journal of Marine Animals and Their Ecology*, 1(7), 29-35.
- Scheidat, M. J., Castro, C., Denkinger, A., González, J., & Adelung, D. (2000). A breeding area for humpback whales (*Megaptera novaeangliae*) off Ecuador. *Journal of Cetacean Research and Management*, 2(3), 165-171.
- Smith, J. N., Grantham, H. S., Gales, N., Double, M. C., Noad, M. J., & Paton, D. (2012). Identification of humpback whale breeding and calving habitat in the Great Barrier Reef. *Marine Ecology Progress Series*, 447, 259-272. <https://doi.org/10.3354/meps09462>
- Stevick, P. T., Aguayo, A., Allen, J., Avila, J. C., Capella, J., Castro, C., . . . Siciliano, S. (2004). Migrations of individually identified humpback whales between the Antarctic Peninsula and South America. *Journal of Cetacean Research and Management*, 6(2), 109-113.
- Stone, G. S., Flórez-González, L., & Katona, S. (1990). Whale migration record. *Nature*, 346, 705. <https://doi.org/10.1038/346705a0>
- Van Waerebeek, K., Djiba, A., Krakstad, J. O., Ould Bilal, A. S., Bamy, I. L., Almeida, A., & Mbye, E. M. (2013). New evidence for a South Atlantic stock of humpback whales wintering on the northwest African continental shelf. *African Zoology*, 48(1), 177-186. <https://doi.org/10.3377/004.048.0120>
- Zerbini, A. N., Andriolo, A., Da Rocha, J. M., Simões-Lopes, P. C., Siciliano, S., Pizzorno, J. L., . . . VanBlaricom, G. R. (2004). Winter distribution and abundance of humpback whales (*Megaptera novaeangliae*) off northeastern Brazil. *Journal of Cetacean Research and Management*, 6(1), 101-107.